

Conjunctivitis

Children can go to school. They will be encouraged to wash their hands to prevent further spread of infection.

Impetigo

Children can go back to school when their lesions are crusted or healed.

Measles, chicken Pox and German Measles

Children should go back to school 5 days after the rash has started. Please let the school know, as pregnant members of staff may be affected.


MUMPS

Children should go back to school 5 days from the start of swollen glands.

Whooping cough

Children should go back to school 5 days after starting antibiotics. Non infectious coughing may continue for many weeks.

Your child...
their future



Medicines in school

Children can come to school even if they are taking medicines, as staff are able to give them their medicine in school. Please make sure the bottle is labelled with your child's name and how often they should have it. Please discuss with the Headteacher.

Should my
child go to
school
today?
th!nk twice

Every day counts



Catch the attendance bug!

Headache, ear ache and stomach ache


Children with headache, ear ache or stomach ache can go to school - just let the staff know they have felt unwell.

Give paracetamol and plenty of fluids to drink.

If headache, ear ache or stomach ache persists.....

EVERY DAY COUNTS

Seek medical advice



HIGH TEMPERATURE

Give paracetamol and plenty to drink. If temperature continues for 3 days or more, seek medical advice.

After paracetamol, if your child feels better, bring them into school.

Coughs and colds

Children should be given paracetamol, plenty of fluids to drink and can be sent to school.

If your child is asthmatic, remember they may need their blue inhaler more often.

Flu and Swine Flu

Children should go back to school when recovered - this is usually about 5 days.

Not Sure ?
Ask For Help.

sore throat,
tonsillitis and
glandular fever

Children should be given paracetamol. Plenty of fluids to drink and can be sent to school.

Diarrhoea and vomiting

Children can return to school 48 hours after the last episode of diarrhoea or 24 hours after the last episode of vomiting.

But do not go swimming for 2 weeks after last episode of diarrhoea.

HEAD LICE

Children can go to school with head lice but they must be treated for the condition to prevent further spread. There is an expectation that parents will be treating their children and family members weekly by wet combing with a nit comb and conditioner.

SEE YOUR SCHOOL NURSE FOR FURTHER ADVICE

Hand, foot and mouth warts and verrucae, athletes foot and molluscum contagioism

Children can go to school. Verrucae should be covered in swimming pools and changing rooms.

Scabies

Children can go back to school after the first treatment. The child and others at home should be treated as well.

THREADWORM

Children can go to school when they have started their treatment.